

DIABETES

Low Blood Glucose (hypoglycemia)

There are many things you can do to manage your blood glucose and live well.

Hypoglycemia is low blood glucose—or low blood sugar—that is below the healthy range. This is usually when your blood glucose is less than 70 mg/dL. You should talk with your diabetes care team about your blood glucose targets, and what level is too low for you.

You may have also heard low blood glucose described as an insulin reaction or insulin shock.

Causes

You might get low blood glucose if you:

- Take certain medicines and eat too few carbohydrates
- Skip or delay meals
- Take too much insulin or diabetes medication (ask your diabetes care team if this applies to you)

Signs and Symptoms

Here's what may happen when your blood glucose is low:

Shaky
Sweaty
Dizzy
Confusion and difficulty speaking

Image: Confusion and difficulty speaking
Image: Confusion and difficulty speaking
Image: Confusion and difficulty speaking

Image: Confusion and difficulty speaking
Image: Confusion and difficulty speaking
Image: Confusion and difficulty speaking

Image: Confusion and difficulty speaking
Image: Confusion and difficulty speaking
Image: Confusion and difficulty speaking

Image: Confusion and difficulty speaking
Image: Confusion and difficulty speaking
Image: Confusion and difficulty speaking

Image: Confusion and difficulty speaking
Image: Confusion and difficulty speaking
Image: Confusion and difficulty speaking

Image: Confusion and difficulty speaking
Image: Confusion and difficulty speaking
Image: Confusion and difficulty speaking

Image: Confusion and difficulty speaking
Image: Confusion and difficulty speaking
Image: Confusion and difficulty speaking

Image: Confusion and difficulty speaking
Image: Confusion and difficulty speaking
Image: Confusion and difficulty speaking

Image: Confusion and difficulty speaking
Image: Confusion and difficulty speaking
Image: Confusion and difficulty speaking

Image: Confusion and difficulty speaking
Image: Confusion and difficulty speaking
Image: Confusion and dimage: Confusion and dimage: Confusion and dimage: Confusion and d

Or you may have no symptoms at all.


If low blood glucose is not treated, it can become severe and may cause you to pass out. If low blood glucose is a problem for you, talk to your doctor or diabetes care team.



- Are more active than usual
- Are sick
- Drink alcohol without eating enough food

Low Blood Glucose (hypoglycemia)

What to do if you think you have low blood glucose



- Check your blood glucose right away if you have any symptoms of low blood glucose
- If you think your blood glucose is low but cannot check it at that time, treat anyway

TREAT

- 4 ounces (½ cup) of regular fruit juice (like orange, apple, or grape juice)
- 4 glucose tablets or 1 tube of glucose gel
- 1 tablespoon of sugar, honey, or corn syrup
- 4 ounces (½ cup) of regular soda pop (not diet)
- 2 tablespoons of raisins



Visit **www.NovoCare.com** for additional resources and helpful information! Point your smartphone camera at the QR code for quick access to the website on your phone.



Novo Nordisk Inc. grants permission to reproduce this piece for nonprofit educational purposes only on condition that the piece is maintained in its original format and that the copyright notice is displayed. Novo Nordisk Inc. reserves the right to revoke this permission at any time.

NovoCare® is a registered trademark of Novo Nordisk A/S.

Novo Nordisk is a registered trademark of Novo Nordisk A/S.

© 2022 Novo Nordisk Printed in the U.S.A. US22PAT00040 October 2022

www.NovoCare.com

