

Getting Started on Once-Weekly OZEMPIC®

OZEMPIC® DOSING SCHEDULE FOR THE PEN THAT DELIVERS 0.25 MG AND 0.5 MG

START



4 WEEKS

The beginning dose is 0.25 mg for the first 4 weeks. This will give your body a chance to get used to the medicine.

- Take Ozempic® once a week, on the same day, exactly as prescribed by your health care provider.
- You can take Ozempic® with or without food.
- You may change the day of the week you use Ozempic® as long as your last dose was given 2 or more days before.
- If you miss a dose of Ozempic®, take the missed dose as soon as possible within 5 days after the missed dose. If more than 5 days have passed, skip the missed dose and take your next dose on the regularly scheduled day.

STAY



WEEK 5

On Week 5, the dose should be increased to 0.5 mg.

My Dose: _____ My Day of Week: Mon Tue Wed Thu Fri Sat Sun

Health Care Provider Notes: _____

Your health care provider should show you how to use Ozempic® before you use it for the first time. Ozempic® is injected under the skin of your stomach (abdomen), thigh, or upper arm. **Do not** inject Ozempic® into a muscle or vein. **Do not** use the same site for each injection.

What is Ozempic®?

Ozempic® (semaglutide) injection 0.5 mg or 1 mg is an injectable prescription medicine used:

- along with diet and exercise to improve blood sugar (glucose) in adults with type 2 diabetes mellitus.
- to reduce the risk of major cardiovascular events such as heart attack, stroke, or death in adults with type 2 diabetes mellitus with known heart disease.

It is not known if Ozempic® can be used in people who have had pancreatitis.

Ozempic® is not for use in people with type 1 diabetes.

It is not known if Ozempic® is safe and effective for use in children under 18 years of age.

Important Safety Information

Do not share your Ozempic® pen with other people, even if the needle has been changed. You may give other people a serious infection, or get a serious infection from them.

Please see Important Safety Information throughout, and Instructions for Use at <https://www.novo-pi.com/ozempic.pdf> for complete instructions. Brief Summary of Information about Ozempic® provided on last page.

ONCE-WEEKLY
OZEMPIC®
semaglutide injection 0.5mg/1mg

Important Safety Information (cont'd)

What is the most important information I should know about Ozempic®?

Ozempic® may cause serious side effects, including:

- **Possible thyroid tumors, including cancer.** Tell your health care provider if you get a lump or swelling in your neck, hoarseness, trouble swallowing, or shortness of breath. These may be symptoms of thyroid cancer. In studies with rodents, Ozempic® and medicines that work like Ozempic® caused thyroid tumors, including thyroid cancer. It is not known if Ozempic® will cause thyroid tumors or a type of thyroid cancer called medullary thyroid carcinoma (MTC) in people.
- Do not use Ozempic® if you or any of your family have ever had MTC, or if you have an endocrine system condition called Multiple Endocrine Neoplasia syndrome type 2 (MEN 2).

Do not use Ozempic® if:

- you or any of your family have ever had MTC or if you have MEN 2.
- you are allergic to semaglutide or any of the ingredients in Ozempic®. See symptoms of serious allergic reaction in "What are the possible side effects of Ozempic®?".

Before using Ozempic®, tell your health care provider if you have any other medical conditions, including if you:

- have or have had problems with your pancreas or kidneys.
- have a history of diabetic retinopathy.
- are pregnant or breastfeeding or plan to become pregnant or breastfeed. It is not known if Ozempic® will harm your unborn baby or passes into your breast milk. You should stop using Ozempic® 2 months before you plan to become pregnant.

Tell your health care provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, herbal supplements, and other medicines to treat diabetes, including insulin or sulfonylureas.

What are the possible side effects of Ozempic®?

Ozempic® may cause serious side effects, including:

- **inflammation of your pancreas (pancreatitis).** Stop using Ozempic® and call your health care provider right away if you have severe pain in your stomach area (abdomen) that will not go away, with or without vomiting. You may feel the pain from your abdomen to your back.
- **changes in vision.** Tell your health care provider if you have changes in vision during treatment with Ozempic®.
- **low blood sugar (hypoglycemia).** Your risk for getting low blood sugar may be higher if you use Ozempic® with another medicine that can cause low blood sugar, such as a sulfonylurea or insulin. **Signs and symptoms of low blood sugar may include:** dizziness or lightheadedness, blurred vision, anxiety, irritability or mood changes, sweating, slurred speech, hunger, confusion or drowsiness, shakiness, weakness, headache, fast heartbeat, and feeling jittery.
- **kidney problems (kidney failure).** In people who have kidney problems, diarrhea, nausea, and vomiting may cause a loss of fluids (dehydration), which may cause kidney problems to get worse. It is important for you to drink fluids to help reduce your chance of dehydration.
- **serious allergic reactions.** Stop using Ozempic® and get medical help right away if you have any symptoms of a serious allergic reaction, including swelling of your face, lips, tongue, or throat; problems breathing or swallowing; severe rash or itching; fainting or feeling dizzy; or very rapid heartbeat.

The most common side effects of Ozempic® may include nausea, vomiting, diarrhea, stomach (abdominal) pain, and constipation.

Please see additional Important Safety Information throughout, and Instructions for Use leaflet at <https://www.novo-pi.com/ozempic.pdf> for complete instructions.

Brief Summary of Information about Ozempic® provided on last page.



Ozempic® also comes in a 1 mg pen for those who need additional blood sugar control.



Ozempic® is a registered trademark of Novo Nordisk A/S.
Novo Nordisk is a registered trademark of Novo Nordisk A/S.
© 2021 Novo Nordisk All rights reserved. US21OZM00485 June 2021

ONCE-WEEKLY
OZEMPIC®
semaglutide injection 0.5mg/1mg

