

You have been prescribed

OZEMPIC®
semaglutide injection 0.5mg|1mg

OH! OZEMPIC® SUPPORT RESOURCES ARE HERE TO HELP



Ozempic® savings

Eligible patients can pay as little as \$25^a for a 1-, 2-, or 3-month prescription.^b Request a card at OzempicSavings.com.



Text-based savings and support

Text^c **BEGIN to 21848** for reminders, tips, and co-pay savings^c via text message.



Talk to your Health Coach

Call **1-866-696-4090**, Monday through Friday, 9 a.m. to 6 p.m. ET, for one-on-one support for using your Ozempic® Pen.



Additional important resources

Visit Ozempic.com/MyJourney for instructional videos, tips, and more from a team of celebrity experts.

^aApplies to eligible, commercially insured patients with coverage for Ozempic®. Month is defined as 28 days. Eligibility and other restrictions apply. Go to SavingsCardEligibility.com for full program details and eligibility requirements.

^bOffer valid for up to 24 months from the date of activation, subject to a maximum savings of \$150 per 1-month, \$300 per 2-month, or \$450 per 3-month prescription.

^cMessage and data rates may apply. Check with your mobile service provider. Message frequency will be based on your selections. Text HELP to 21848 for help. Text STOP to 21848 to quit. See Terms and Conditions of Use at Ozempic.com.

Open your phone camera and scan this code or visit Ozempic.com to access **Prescribing Information and Medication Guide**.

Find more tools and resources on OzempicQuickResources.com

For the pharmacist: For any questions on how to process the Ozempic® savings offer, please call **1-800-433-4893**.



Please see next page for brief summary of important information about Ozempic®.

Ozempic® is a registered trademark of Novo Nordisk A/S.
Novo Nordisk is a registered trademark of Novo Nordisk A/S.
© 2021 Novo Nordisk All rights reserved. US21OZM00924 December 2021

OZEMPIC®
semaglutide injection 0.5mg|1mg

Brief Summary of information about OZEMPIC® (semaglutide) injection	How should I use OZEMPIC®?
<div>  <div> Rx Only This information is not comprehensive. <ul style="list-style-type: none"> Talk to your healthcare provider or pharmacist Visit www.novo-pi.com/ozempic.pdf to obtain the FDA-approved product labeling Call 1-888-693-6742 </div> </div>	<ul style="list-style-type: none"> OZEMPIC® is injected under the skin (subcutaneously) of your stomach (abdomen), thigh, or upper arm. Do not inject OZEMPIC® into a muscle (intramuscularly) or vein (intravenously). Do not mix insulin and OZEMPIC® together in the same injection. Change (rotate) your injection site with each injection. Do not use the same site for each injection. Talk to your healthcare provider about how to prevent, recognize and manage low blood sugar (hypoglycemia), high blood sugar (hyperglycemia), and problems you have because of your diabetes.
<p>Do not share your OZEMPIC® pen with other people, even if the needle has been changed. You may give other people a serious infection, or get a serious infection from them.</p>	<p>What are the possible side effects of OZEMPIC®? OZEMPIC® may cause serious side effects, including:</p>
<p>Read this Medication Guide before you start using OZEMPIC® and each time you get a refill. There may be new information. This information does not take the place of talking to your healthcare provider about your medical condition or your treatment.</p>	<p>See “What is the most important information I should know about OZEMPIC®?”</p>
<p>What is the most important information I should know about OZEMPIC®? OZEMPIC® may cause serious side effects, including:</p> <ul style="list-style-type: none"> Possible thyroid tumors, including cancer. Tell your healthcare provider if you get a lump or swelling in your neck, hoarseness, trouble swallowing, or shortness of breath. These may be symptoms of thyroid cancer. In studies with rodents, OZEMPIC® and medicines that work like OZEMPIC® caused thyroid tumors, including thyroid cancer. It is not known if OZEMPIC® will cause thyroid tumors or a type of thyroid cancer called medullary thyroid carcinoma (MTC) in people. Do not use OZEMPIC® if you or any of your family have ever had a type of thyroid cancer called medullary thyroid carcinoma (MTC), or if you have an endocrine system condition called Multiple Endocrine Neoplasia syndrome type 2 (MEN 2). 	<ul style="list-style-type: none"> inflammation of your pancreas (pancreatitis). Stop using OZEMPIC® and call your healthcare provider right away if you have severe pain in your stomach area (abdomen) that will not go away, with or without vomiting. You may feel the pain from your abdomen to your back. changes in vision. Tell your healthcare provider if you have changes in vision during treatment with OZEMPIC®. low blood sugar (hypoglycemia). Your risk for getting low blood sugar may be higher if you use OZEMPIC® with another medicine that can cause low blood sugar, such as a sulfonylurea or insulin. Signs and symptoms of low blood sugar may include: <ul style="list-style-type: none"> dizziness or light-headedness blurred vision anxiety, irritability, or mood changes sweating slurred speech hunger confusion or drowsiness shakiness weakness headache fast heartbeat feeling jittery kidney problems (kidney failure). In people who have kidney problems, diarrhea, nausea, and vomiting may cause a loss of fluids (dehydration) which may cause kidney problems to get worse. It is important for you to drink fluids to help reduce your chance of dehydration. serious allergic reactions. Stop using OZEMPIC® and get medical help right away, if you have any symptoms of a serious allergic reaction including: <ul style="list-style-type: none"> swelling of your face, lips, tongue or throat problems breathing or swallowing severe rash or itching fainting or feeling dizzy very rapid heartbeat
<p>What is OZEMPIC®? OZEMPIC® is an injectable prescription medicine used:</p> <ul style="list-style-type: none"> along with diet and exercise to improve blood sugar (glucose) in adults with type 2 diabetes mellitus. to reduce the risk of major cardiovascular events such as heart attack, stroke or death in adults with type 2 diabetes mellitus with known heart disease. <p>It is not known if OZEMPIC® can be used in people who have had pancreatitis. OZEMPIC® is not for use in people with type 1 diabetes. It is not known if OZEMPIC® is safe and effective for use in children under 18 years of age.</p>	<p>The most common side effects of OZEMPIC® may include nausea, vomiting, diarrhea, stomach (abdominal) pain and constipation.</p> <p>Talk to your healthcare provider about any side effect that bothers you or does not go away. These are not all the possible side effects of OZEMPIC®.</p> <p>Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.</p>
<p>Do not use OZEMPIC® if:</p> <ul style="list-style-type: none"> you or any of your family have ever had a type of thyroid cancer called medullary thyroid carcinoma (MTC) or if you have an endocrine system condition called Multiple Endocrine Neoplasia syndrome type 2 (MEN 2). you have had a serious allergic reaction to semaglutide or any of the ingredients in OZEMPIC®. Symptoms of a serious allergic reaction include: <ul style="list-style-type: none"> swelling of your face, lips, tongue or throat problems breathing or swallowing severe rash or itching fainting or feeling dizzy very rapid heartbeat 	
<p>Before using OZEMPIC®, tell your healthcare provider if you have any other medical conditions, including if you:</p> <ul style="list-style-type: none"> have or have had problems with your pancreas or kidneys. have a history of diabetic retinopathy. are pregnant or plan to become pregnant. It is not known if OZEMPIC® will harm your unborn baby. You should stop using OZEMPIC® 2 months before you plan to become pregnant. Talk to your healthcare provider about the best way to control your blood sugar if you plan to become pregnant or while you are pregnant. are breastfeeding or plan to breastfeed. It is not known if OZEMPIC® passes into your breast milk. You should talk with your healthcare provider about the best way to feed your baby while using OZEMPIC®. 	
<p>Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. OZEMPIC® may affect the way some medicines work and some medicines may affect the way OZEMPIC® works.</p>	
<p>Before using OZEMPIC®, talk to your healthcare provider about low blood sugar and how to manage it. Tell your healthcare provider if you are taking other medicines to treat diabetes, including insulin or sulfonylureas. Know the medicines you take. Keep a list of them to show your healthcare provider and pharmacist when you get a new medicine.</p>	