

# Checking your blood sugar

## How frequent blood sugar checks can help you

Checking your blood sugar yourself is an important part of managing diabetes. Checking often will tell you:

- If your insulin or other diabetes medicine is working
- How physical activity and the foods you eat affect your blood sugar

You'll usually feel better and have more energy when your blood sugar stays at or near normal. Managing your blood sugar can also reduce your risk of developing problems from diabetes.



## How to check your blood sugar

You can check your own blood sugar using a meter. Many different kinds of blood sugar meters are available today. Your diabetes care team can help you choose one and show you how to use it.

## When to check your blood sugar

You and your diabetes care team will decide when and how often you will check your blood sugar. The table below shows some times when you might want to check and why.

When your team may want you to check	Why you should check
When you wake up	To see if your blood sugar is staying under control while you're asleep (called fasting blood sugar)
Before meals	To know what your blood sugar is before you eat and to know if you will need to adjust your mealtime (prandial) insulin
1 or 2 hours after you start your meal	To see how the food you eat and insulin dose affect your blood sugar
Before, during, and after physical activity (depending on the length of time of your activity)	To see how being active affects your blood sugar
At bedtime	Depending on the medicine that you take

## Keeping a blood sugar tracker

It's important to write down your blood sugar levels so that you can keep track of what makes them go up or down.

See the last page for a tracker that you can use to record your blood sugar and write down other important information for your diabetes care team.



**For mobile tracking, try the Cornerstones4Care® Powered by Glooko app!**

## Checking your blood sugar

Time	Goals for many nonpregnant adults with diabetes	Your goals
Before meals	80 to 130 mg/dL	_____
1 to 2 hours after the start of a meal	Less than 180 mg/dL	_____
A1C	Less than 7%	_____

American Diabetes Association. Standards of medical care in diabetes—2019. *Diabetes Care*. 2019;42(suppl 1):S1-S193.


### Setting your blood sugar goals

The table above lists blood sugar goals for many adults with diabetes. You and your diabetes care team will set the goals that are right for you. Write your goals in the last column.

### Knowing your A1C

The A1C test measures your estimated average blood sugar level over approximately 3 months. It's like a "memory" of your blood sugar levels. It shows how well you're controlling your blood sugar levels over time. Your A1C and your blood sugar levels go up and down together. The table to the right shows how A1C relates to the estimated average blood sugar level.

Lowering your A1C to below 7% reduces your risk of problems from diabetes. Therefore, the A1C goal for most people is less than 7%.



It is recommended that you get an A1C test:

- At least 2 times a year if your blood sugar is under good control
- 4 times a year if you are not meeting your goals or if your treatment has changed

A1C level	Average blood sugar
6%	126 mg/dL
7%	154 mg/dL
8%	183 mg/dL
9%	212 mg/dL
10%	240 mg/dL
11%	269 mg/dL
12%	298 mg/dL

American Diabetes Association. Standards of medical care in diabetes—2019. *Diabetes Care*. 2019;42(suppl 1):S1-S193.

If you have any questions, be sure to talk with your diabetes care team. They are there to help!

For more information, visit [Cornerstones4Care.com](http://Cornerstones4Care.com)



Novo Nordisk Inc. grants permission to reproduce this piece for nonprofit educational purposes only on condition that the piece is maintained in its original format and that the copyright notice is displayed. Novo Nordisk Inc. reserves the right to revoke this permission at any time.

Cornerstones4Care® is a registered trademark of Novo Nordisk A/S.

Novo Nordisk is a registered trademark of Novo Nordisk A/S.

© 2019 Novo Nordisk All rights reserved. US19DI00085 June 2019 [Cornerstones4Care.com](http://Cornerstones4Care.com)

## How to use your blood sugar tracker

Cornerstones4Care®

### Your blood sugar tracker

Diabetes changes over time. That is why your treatment may also need to change over time. For example, your doctor may tell you to add a basal insulin to your diabetes pills. Or if you already take a basal insulin, your doctor may tell you to add mealtime insulin.

The blood sugar tracker on the next page can help you keep track of your diabetes medicines and any changes in dosage and timing that your doctor may tell you to make. It can also help you keep track of how much mealtime insulin you took and when you took it.

Your diabetes care team can provide you with a blood sugar tracker and show you how to use it. Or call **1-800-727-6500** to have a tracker sent to you.

### Here's how to use the blood sugar tracker

- 1 Write down the date for the start of the week. (You can start tracking on any day of the week)
- 2 Write the name(s) and dose(s) of your non-insulin diabetes medicine(s), such as pills or non-insulin injectable medicines
- 3 Write the time and your blood sugar readings in the "before" and "after" spaces. After-meal readings are usually taken 1 to 2 hours after you start your meal. Nighttime readings may be taken during the night as needed
- 4 If you are counting carbs, write how many grams of carbs you ate
- 5 If your doctor has told you to use mealtime insulin when you eat, write your dose here
- 6 If your doctor has told you to use long-acting insulin, write your dose and time(s) here. Long-acting insulin is taken either once or twice a day
- 7 Add notes on anything else you might want to track (such as blood pressure, activity, or weight)

### EXAMPLE

**Your blood sugar tracker** 1 Date: 4 / 22 / 2019

		BLOOD SUGAR RESULTS*									
		Breakfast		Lunch		Dinner		Bedtime	Night		
		BEFORE	AFTER	BEFORE	AFTER	BEFORE	AFTER	Time			Time
2	<b>FRIDAY</b>										
	Non-insulin medicine and dose	10 mg medication A									
		Time	Time	Time	Time	Time	Time				
		6 am	8 am		1:30 pm	6:15 pm		11 pm			3:30 am
					148	91		90			140
		mg/dL	mg/dL	mg/dL	mg/dL	mg/dL	mg/dL	mg/dL			mg/dL
4	Carb intake	18 grams		21 grams		26 grams		2 grams			
5	Mealtime insulin dose	<input type="checkbox"/> Breakfast _____ units		<input type="checkbox"/> Lunch _____ units		<input checked="" type="checkbox"/> Dinner 2 units		Other			
6	Long-acting insulin dose	90 units 150 time		If needed at dinner or bedtime _____ units _____ time				BP: 120/80			
	<b>SATURDAY</b>										
	Non-insulin medicine and dose	10 mg medication A									
		Time	Time	Time	Time	Time	Time				
		6 am	8 am								
		90	150								
		mg/dL	mg/dL	mg/dL	mg/dL	mg/dL	mg/dL	mg/dL			mg
	Carb intake	20 grams									
	Mealtime insulin dose	<input type="checkbox"/> Breakfast _____ units		<input type="checkbox"/> Lunch _____ units		<input checked="" type="checkbox"/> Dinner 2 units		Other			
	Long-acting insulin dose	_____ units _____ time		If needed at dinner or bedtime _____ units _____ time				30 minute walk			

\*You and your diabetes care team will decide the best times for you to check your blood sugar.

# Your blood sugar tracker

Date: \_\_\_\_ / \_\_\_\_ / \_\_\_\_

		BLOOD SUGAR RESULTS*									
		Breakfast		Lunch		Dinner		Bedtime	Night		
Non-insulin medicine and dose	<b>MONDAY</b>	BEFORE	AFTER	BEFORE	AFTER	BEFORE	AFTER	Time	Time		
		Time	Time	Time	Time	Time	Time				
		mg/dL	mg/dL	mg/dL	mg/dL	mg/dL	mg/dL		mg/dL		
		Carb intake ▶									
		Mealtime insulin dose ▶ <input type="checkbox"/> Breakfast _____ units		<input type="checkbox"/> Lunch _____ units		<input type="checkbox"/> Dinner _____ units		Other ▶			
	Long-acting insulin dose ▶ _____ units _____ time		If needed at dinner or bedtime		_____ units _____ time						
Non-insulin medicine and dose	<b>TUESDAY</b>	BEFORE	AFTER	BEFORE	AFTER	BEFORE	AFTER	Time	Time		
		Time	Time	Time	Time	Time	Time				
		mg/dL	mg/dL	mg/dL	mg/dL	mg/dL	mg/dL		mg/dL		
		Carb intake ▶									
		Mealtime insulin dose ▶ <input type="checkbox"/> Breakfast _____ units		<input type="checkbox"/> Lunch _____ units		<input type="checkbox"/> Dinner _____ units		Other ▶			
	Long-acting insulin dose ▶ _____ units _____ time		If needed at dinner or bedtime		_____ units _____ time						
Non-insulin medicine and dose	<b>WEDNESDAY</b>	BEFORE	AFTER	BEFORE	AFTER	BEFORE	AFTER	Time	Time		
		Time	Time	Time	Time	Time	Time				
		mg/dL	mg/dL	mg/dL	mg/dL	mg/dL	mg/dL		mg/dL		
		Carb intake ▶									
		Mealtime insulin dose ▶ <input type="checkbox"/> Breakfast _____ units		<input type="checkbox"/> Lunch _____ units		<input type="checkbox"/> Dinner _____ units		Other ▶			
	Long-acting insulin dose ▶ _____ units _____ time		If needed at dinner or bedtime		_____ units _____ time						
Non-insulin medicine and dose	<b>THURSDAY</b>	BEFORE	AFTER	BEFORE	AFTER	BEFORE	AFTER	Time	Time		
		Time	Time	Time	Time	Time	Time				
		mg/dL	mg/dL	mg/dL	mg/dL	mg/dL	mg/dL		mg/dL		
		Carb intake ▶									
		Mealtime insulin dose ▶ <input type="checkbox"/> Breakfast _____ units		<input type="checkbox"/> Lunch _____ units		<input type="checkbox"/> Dinner _____ units		Other ▶			
	Long-acting insulin dose ▶ _____ units _____ time		If needed at dinner or bedtime		_____ units _____ time						
Non-insulin medicine and dose	<b>FRIDAY</b>	BEFORE	AFTER	BEFORE	AFTER	BEFORE	AFTER	Time	Time		
		Time	Time	Time	Time	Time	Time				
		mg/dL	mg/dL	mg/dL	mg/dL	mg/dL	mg/dL		mg/dL		
		Carb intake ▶									
		Mealtime insulin dose ▶ <input type="checkbox"/> Breakfast _____ units		<input type="checkbox"/> Lunch _____ units		<input type="checkbox"/> Dinner _____ units		Other ▶			
	Long-acting insulin dose ▶ _____ units _____ time		If needed at dinner or bedtime		_____ units _____ time						
Non-insulin medicine and dose	<b>SATURDAY</b>	BEFORE	AFTER	BEFORE	AFTER	BEFORE	AFTER	Time	Time		
		Time	Time	Time	Time	Time	Time				
		mg/dL	mg/dL	mg/dL	mg/dL	mg/dL	mg/dL		mg/dL		
		Carb intake ▶									
		Mealtime insulin dose ▶ <input type="checkbox"/> Breakfast _____ units		<input type="checkbox"/> Lunch _____ units		<input type="checkbox"/> Dinner _____ units		Other ▶			
	Long-acting insulin dose ▶ _____ units _____ time		If needed at dinner or bedtime		_____ units _____ time						
Non-insulin medicine and dose	<b>SUNDAY</b>	BEFORE	AFTER	BEFORE	AFTER	BEFORE	AFTER	Time	Time		
		Time	Time	Time	Time	Time	Time				
		mg/dL	mg/dL	mg/dL	mg/dL	mg/dL	mg/dL		mg/dL		
		Carb intake ▶									
		Mealtime insulin dose ▶ <input type="checkbox"/> Breakfast _____ units		<input type="checkbox"/> Lunch _____ units		<input type="checkbox"/> Dinner _____ units		Other ▶			
	Long-acting insulin dose ▶ _____ units _____ time		If needed at dinner or bedtime		_____ units _____ time						

If you'd like to use an online blood sugar tracker, you can find one at [Cornerstones4Care.com](http://Cornerstones4Care.com)

\*You and your diabetes care team will decide the best times for you to check your blood sugar.

