



RYBELSUS[®]
semaglutide tablets 7mg | 14mg

Your Guide to Taking RYBELSUS[®]

Pay as little as \$10 for up
to a 3-month prescription.^a
Text^b **READY** to 21848.

^aEligible, commercially insured patients only. For 3 mg dose, pay as little as \$10 each month. Restrictions apply.

^bMessage and data rates may apply. Check with your mobile service provider. See Terms and Conditions of Use at RYBELSUS.com.

Please see Important Safety Information throughout, and [click here](#) for Prescribing Information and Medication Guide.

Getting Started on RYBELSUS®

START ON 3 mg




The beginning dose is 3 mg for the first 30 days. This will give your body a chance to get used to the medicine.




STAY ON 7 mg



When you finish all of the 3 mg tablets, start your 7 mg prescription.

 Take RYBELSUS® exactly as your healthcare provider tells you to ensure that it works as planned to lower your blood sugar.

 Remember to pick up your prescription from your pharmacy.

Why does my RYBELSUS® dose change?

The beginning dose is 3 mg for the first 30 days to help your body get used to the medicine.

Your healthcare provider prescribed a 7 mg dose to help lower your blood sugar.

If more blood sugar control is needed, your healthcare provider may prescribe the 14 mg dose.

 Please keep reading to learn more about starting RYBELSUS®.

What is RYBELSUS®?

RYBELSUS® (semaglutide) tablets 7 mg or 14 mg is a prescription medicine used along with diet and exercise to improve blood sugar (glucose) in adults with type 2 diabetes.

- RYBELSUS® is not recommended as the first choice of medicine for treating diabetes
- It is not known if RYBELSUS® can be used in people who have had pancreatitis
- RYBELSUS® is not for use in people with type 1 diabetes
- It is not known if RYBELSUS® is safe and effective for use in children under 18 years of age

Please see Important Safety Information throughout, and [click here](#) for Prescribing Information and Medication Guide.

RYBELSUS®
semaglutide tablets 7mg | 14mg

Take RYBELSUS® the Same Way, Every Day

For RYBELSUS® to work as planned to lower blood sugar:



Take on an empty stomach
when you first wake up.

An empty stomach means no food,
drink, oral medications, vitamins,
or supplements.



Take with no more than
4 ounces of water.

Take RYBELSUS® with a sip of plain
water, no more than 4 ounces.



Wait 30 minutes before eating,
drinking, or taking other oral medications.

After 30 minutes, you can eat, drink, or
take other oral medications,
vitamins, or supplements.

Find answers to frequently asked
questions at [RYBELSUSFAQs.com](https://www.rybelsus.com/FAQs).

💡 If you miss a dose, the missed dose should
be skipped, and the next dose should be
taken the following day.

Please see Important Safety Information throughout,
and [click here](#) for Prescribing Information and Medication Guide.

RYBELSUS®
semaglutide tablets 7mg | 14mg

How to Store RYBELSUS®

For RYBELSUS® to work as planned to lower blood sugar:



Store tablets in the closed RYBELSUS® bottle until you are ready to take one.




Do not use a pill organizer or other container to store RYBELSUS® tablets.



Swallow tablet whole. Do not cut, crush, or chew.

 Store at room temperature 68°F to 77°F.

 RYBELSUS® comes in a child-resistant bottle. Keep out of reach of children.

Keeping the Blue Cap Tight Helps the Pills Work Right



The blue cap on the RYBELSUS® bottle is specially designed to keep your tablets protected so they can work as planned to lower your blood sugar. It is important that your RYBELSUS® tablets stay in the bottle with the cap on until you are ready to take one.

Important Safety Information

What is the most important information I should know about RYBELSUS®?

RYBELSUS® may cause serious side effects, including:

- **Possible thyroid tumors, including cancer.** Tell your healthcare provider if you get a lump or swelling in your neck, hoarseness, trouble swallowing, or shortness of breath. These may be symptoms of thyroid cancer. In studies with rodents, RYBELSUS® and medicines that work like RYBELSUS® caused thyroid tumors, including thyroid cancer. It is not known if RYBELSUS® will cause thyroid tumors or a type of thyroid cancer called medullary thyroid carcinoma (MTC) in people

Please see Important Safety Information throughout, and [click here](#) for Prescribing Information and Medication Guide.

RYBELSUS®
semaglutide tablets 7mg | 14mg

Common Side Effects

The most common side effects with RYBELSUS® may include nausea, stomach (abdominal) pain, diarrhea, decreased appetite, vomiting, and constipation. Nausea, vomiting, and diarrhea are most common when starting RYBELSUS®.^a

If you experience nausea while taking RYBELSUS®, please talk to your healthcare provider.

Here are some nausea tips that may be helpful:

- ✓ Slowly eat smaller, more frequent meals
- ✓ Eat foods that are light and bland, like saltine crackers or plain bread
- ✓ Avoid fried, greasy, or sweet foods
- ✓ Drink clear or ice-cold drinks

^aIn studies, GI side effects were more common in people taking RYBELSUS® than in people taking placebo, and people taking RYBELSUS® were more likely to stop treatment because of these side effects. Nausea, vomiting, and diarrhea may cause a loss of fluids (dehydration), which could cause existing kidney problems to get worse (including kidney failure).

Important Safety Information

Do not use RYBELSUS® if:

- you or any of your family have ever had MTC, or if you have an endocrine system condition called Multiple Endocrine Neoplasia syndrome type 2 (MEN 2)
- you have had a serious allergic reaction to semaglutide or any of the ingredients in RYBELSUS®. See symptoms of serious allergic reaction in **“What are the possible side effects of RYBELSUS®?”**

Before using RYBELSUS®, tell your healthcare provider if you have any other medical conditions, including if you:

- have or have had problems with your pancreas or kidneys
- have a history of vision problems related to your diabetes
- are pregnant or plan to become pregnant. It is not known if RYBELSUS® will harm your unborn baby. You should stop using RYBELSUS® 2 months before you plan to become pregnant. Talk to your healthcare provider about the best way to control your blood sugar if you plan to become pregnant or while you are pregnant
- are breastfeeding or plan to breastfeed. Breastfeeding is not recommended during treatment with RYBELSUS®

Please see Important Safety Information throughout, and [click here](#) for Prescribing Information and Medication Guide.

RYBELSUS®
semaglutide tablets 7mg | 14mg

RYBELSUS® Savings



SAVE ON RYBELSUS®



FOR UP
TO A



MONTH
PRESCRIPTION

^aTo receive offer, 7 mg or 14 mg dose prescription must be for a 1-, 2-, or 3-month supply. For 3 mg dose, pay as little as \$10 each month. Eligible, commercially insured patients only. Maximum savings of \$300 per 1-month supply, \$600 per 2-month supply, or \$900 per 3-month supply. Eligibility and other restrictions apply. See Terms and Conditions of Use at RYBELSUS.com.

Who qualifies for the RYBELSUS® Savings Offer?

To qualify, commercial insurance coverage is required. Eligibility and other restrictions apply. Visit [SaveOnR.com](https://www.saveonr.com) for full program details and eligibility requirements.

Text or go online to get a Savings Offer.

Sign Up for RYBELSUS® Support



RYBELSUS® SUPPORT

Sign up for text messages with pill reminders, helpful information, and motivational messages!

Text^a **READY** to **21848** to get started.

^aMessage and data rates may apply. Check with your mobile service provider. See Terms and Conditions of Use at RYBELSUS.com.



ONE-ON-ONE, LIVE SUPPORT

There's more support from a Diabetes Health Coach, if you need it.

Call **1-833-275-2233**

Monday through Friday,
9:00 AM – 6:00 PM ET.

Please see Important Safety Information throughout, and [click here](#) for Prescribing Information and Medication Guide.


RYBELSUS®
semaglutide tablets 7mg | 14mg

Track Your Dose

Print the image of the bottle below to track your first 30 days on the 3 mg dose of RYBELSUS®. Check off one of the spaces below after you take your daily tablet. Building this habit may help you fit RYBELSUS® into your wake-up routine.

START ON 3 mg

1	2	3	4	5
6	7	8	9	10
11	12	13	14	15
16	17	18	19	20
21	22	23	24	25
26	27	28	29	30

 Keep your RYBELSUS® tablets in the RYBELSUS® bottle with the blue cap on to ensure that they can work as planned to lower your blood sugar.

Please see Important Safety Information throughout, and [click here](#) for Prescribing Information and Medication Guide.

RYBELSUS®
semaglutide tablets 7mg | 14mg

Talking to Your Healthcare Provider

For your next visit, it may be helpful to print this page and write down your questions to help guide your conversation with your healthcare provider.

Example: What will my next dosage be?



Scan this QR code for helpful information about RYBELSUS[®].

Please see Important Safety Information throughout, and [click here](#) for Prescribing Information and Medication Guide.



Important Safety Information (cont'd)

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. RYBELSUS® may affect the way some medicines work and some medicines may affect the way RYBELSUS® works.

What are the possible side effects of RYBELSUS®?

RYBELSUS® may cause serious side effects, including:

- **inflammation of your pancreas (pancreatitis).** Stop using RYBELSUS® and call your healthcare provider right away if you have severe pain in your stomach area (abdomen) that will not go away, with or without vomiting. You may feel the pain from your abdomen to your back
- **changes in vision.** Tell your healthcare provider if you have changes in vision during treatment with RYBELSUS®
- **low blood sugar (hypoglycemia).** Your risk for getting low blood sugar may be higher if you use RYBELSUS® with another medicine that can cause low blood sugar, such as a sulfonylurea or insulin. **Signs and symptoms of low blood sugar may include:** dizziness or lightheadedness, blurred vision, anxiety, irritability or mood changes, sweating, slurred speech, hunger, confusion or drowsiness, shakiness, weakness, headache, fast heartbeat, and feeling jittery
- **kidney problems (kidney failure).** In people who have kidney problems, diarrhea, nausea, and vomiting may cause a loss of fluids (dehydration), which may cause kidney problems to get worse. It is important for you to drink fluids to help reduce your chance of dehydration
- **serious allergic reactions.** Stop using RYBELSUS® and get medical help right away, if you have any symptoms of a serious allergic reaction, including swelling of your face, lips, tongue, or throat; problems breathing or swallowing; severe rash or itching; fainting or feeling dizzy; or very rapid heartbeat
- **gallbladder problems.** Gallbladder problems have happened in some people who take RYBELSUS®. Tell your healthcare provider right away if you get symptoms of gallbladder problems, which may include: pain in your upper stomach (abdomen), yellowing of skin or eyes (jaundice), fever, and clay-colored stools

The most common side effects of RYBELSUS® may include nausea, stomach (abdominal) pain, diarrhea, decreased appetite, vomiting, and constipation. Nausea, vomiting, and diarrhea are most common when you first start RYBELSUS®.

Please see Important Safety Information throughout and [click here](#) for Prescribing Information and Medication Guide.

RYBELSUS®
semaglutide tablets 7mg | 14mg

Savings and support are just a text away.

Text^a **READY** to **21848** to get started.

^aMessage and data rates may apply. Check with your mobile service provider. Message frequency will be based on your selections. Text HELP to 21848 for help. Text STOP to 21848 to quit. See Terms and Conditions of Use at RYBELSUS.com.

HELPFUL TIPS

- ✓ Track your blood sugar and share with your healthcare provider during your next visit
- ✓ Make activity and healthy eating part of your daily routine
- ✓ Remember to schedule all your appointments, like eye and foot checks
- ✓ Forming a new habit is easier when it fits into your daily routine, like brushing your teeth

Call 1-833-275-2233 or visit RYBELSUS.com for more information.

Please see Important Safety Information throughout and [click here](#) for Prescribing Information and Medication Guide.

RYBELSUS® is a registered trademark of Novo Nordisk A/S.

Novo Nordisk is a registered trademark of Novo Nordisk A/S.

© 2022 Novo Nordisk All rights reserved. US21RYB00824 June 2022



RYBELSUS®
semaglutide tablets 7mg | 14mg